

DTP/00/85
gr-qc/0009102

Cosmic strings in axionic-dilatonic gravity.

Caroline Santos

Departamento de Física da Faculdade de Ciências da Universidade do Porto, Rua do
Campo Alegre 687, 4150-Porto, Portugal
Centre for Particle Theory, Department of Mathematical Sciences, South Road,
Durham, DH1 3LE.

Abstract

We first consider local cosmic strings in dilaton-axion gravity and show that they are singular solutions. Then we take a supermassive Higgs limit and present expressions for the fields at far distances from the core by applying a Pecci-Quinn and a duality transformation to the dilatonic Melvin's magnetic universe.

PACS numbers: 04.70.-b, 11.27.+d

1 Introduction

Topological defects of various kinds are predicted to have occurred during phase transitions in the early universe. Depending on the nature of the phase transition and the configuration of the fields involved in spontaneous symmetry-breaking, topological defects can be three, two or one dimensional, such as textures, domain walls or strings, and also pointlike such as monopoles. Their cosmological implications have been quite well explored in the context of general relativity (for a review see [1]). In particular the cosmic strings which have attracted the most attention are those produced at GUT scales. Such strings generate large angle cosmic microwave background anisotropies of roughly the observed order of magnitude [2]. Their gravitational effects are weak and their static spacetimes are non-singular [3, 4].

On the other hand, at sufficiently high energy scales the Einstein's theory of general relativity which is extremely successful at describing the dynamics of our solar system, and indeed the observable universe, probably does not describe gravity accurately [5] as first postulated by Dirac [6].

Partially motivated by the Dirac's idea and the possible existence of extra dimensions of the spacetime proposed by Kaluza and Klein [7], Jordan, Brans and Dicke (JBD) [8] proposed a theory of gravity whose purpose was to incorporate Mach's Principle. In this way the variations on the inertial mass of a body caused by the surrounding Universe, assumed in that Principle, could be justified from the variations of the gravitational constant. Even so the Principle is not consistently justified, as it does not explain gravity in vacuum.

The implications of the JBD action on cosmic string models have been well explored [9, 10]. In particular the case of the low-energy string action was explored in [11] which showed that in the presence of a dilaton the gravitational field of cosmic strings is surprisingly close to that of an Einstein cosmic string on cosmological distance scales. For a massless dilaton it was shown that there are long range dilatonic effects on the very large scale and the spacetime is not asymptotically locally flat in the Einstein and string frames. On scales of cosmological interest, however, the metric is conical in the Einstein frame, and conformally conical in the string frame. Meanwhile for a massive dilaton and apart from distances of order the Compton wavelength of the dilaton, the

long-range structure of the string is as for Einstein gravity, and so the metric asymptotes a conical metric, in both string and Einstein frames. The main exception to these results is for the special value where the coupling of the dilaton to the vortex fields is canonical, i.e. the vortex fields couple to the metric in the string frame, for which there are no long range effects (other than the deficit angle) and the dilaton merely shifts the value of the dilaton between the core and infinity by a constant. For a critical Higgs coupling this constant vanishes and there is no effect at all on the dilaton. The dilatonic string is then the same as the Einstein one.

Meanwhile, low energy string gravity also generically contains another field - the axion - which could in principle also couple to the cosmic string. Of course, in the cosmic super string solution of Dabholkar et al. [12] the axion couples to the string. However here we wish to consider the simple question of whether and how, at the low energy level, the presence of the axion effects the field of a local vortex solution to some field theory, using the $U(1)$ abelian Higgs theory as a model.

Therefore the layout of this paper is as follows: Section 2 shows the singular behaviour of the axionic-dilatonic local strings by examining the gravi-axion-dilaton field. Section 3 presents a far field solution by taking a supermassive Higgs limit. Finally we conclude.

2 Singular axionic-dilatonic strings

We are interested in the behaviour of a static self gravitating string whose metric is given by

$$ds^2 = e^\gamma \left[dt^2 - dR^2 - dz^2 \right] - \alpha^2 e^{-\gamma} d\varphi^2 . \quad (1)$$

when the gravitational interactions take a form typical of low energy string theory [13].

We take an empirical approach to cosmic strings in this background theory, not concerning ourselves with the origin of the fields that form the vortex, but inputting “*by hand*” the abelian-Higgs lagrangian given by

$$\mathcal{L}[\Phi, A_c] = D_a \Phi^\dagger D^a \Phi - \frac{1}{4} \tilde{F}_{ab} \tilde{F}^{ab} - \frac{\chi}{4} (\Phi^\dagger \Phi - \eta^2)^2 \quad (2)$$

where $D_c = \nabla_c + ieA_c$ is the usual gauge covariant derivative, $\tilde{F}_{ab}$ is the field strength of the gauge field A_c , η is the symmetry breaking scale and χ is a parameter related with

the energy density of the string, μ , by

$$\mu \sim \frac{1}{\sqrt{\chi}\eta} . \quad (3)$$

To take account of the (unknown) coupling of the cosmic string to the dilaton ϕ , we consider a reasonably general form for the interaction of the vortex with the dilaton assuming that the Abelian-Higgs lagrangian couples to the dilaton via an arbitrary coupling, $e^{2a\phi} \mathcal{L}$, in the string frame as in the action:

$$\hat{S} = \int d^4x \sqrt{-\hat{g}} \left[e^{-2\phi} \left(-\hat{R} - 4(\hat{\nabla}\phi)^2 - \hat{V}(\phi) + \frac{1}{12} \hat{H}_{\mu\nu\lambda}^2 \right) + e^{2a\phi} \hat{\mathcal{L}} \right] \quad (4)$$

where $\hat{V}(\phi)$ is the dilaton potential and the axion field is described by an antisymmetric tensor $B_{\nu\lambda}$ with antisymmetric field strength $\hat{H}_{\mu\nu\lambda}$ given by: $\hat{H}_{\mu\nu\lambda} = \partial_{[\mu} B_{\nu\lambda]}$.

We now write instead the action in terms of the “*Einstein*” metric, defined as

$$g_{ab} = e^{-2\phi} \hat{g}_{ab} \quad (5)$$

in which the gravitational part of the action appears in the more familiar Einstein form:

$$S = \int d^4x \sqrt{-g} \left[-R + 2(\nabla\phi)^2 - V(\phi) + \frac{1}{12} e^{-4\phi} H_{\mu\nu\lambda}^2 + 2\epsilon e^{2(a+2)\phi} \mathcal{L}\{X, P, e^{2\phi}g\} \right] \quad (6)$$

and write the “*Einstein*” equations to get

$$G_{ab} = \epsilon e^{2(a+2)\phi} T_{ab} + S_{ab} + N_{ab} \quad (7)$$

with T_{ab} , S_{ab} and N_{ab} the energy-momentum tensors for the string, the dilaton and the axion given respectively by:

$$T_{ab} = 2 \frac{\delta \mathcal{L}[X, P, e^{2\phi}g]}{\delta g^{ab}} = 2e^{-2\phi} [\nabla_a X \nabla_b X + X^2 P_a P_b] - 2\beta e^{-4\phi} F_{ac} F_b^c - \mathcal{L} g_{ab} , \quad (8)$$

$$S_{ab} = 2\nabla_a \phi \nabla_b \phi + \frac{1}{2} V(\phi) g_{ab} - (\nabla\phi)^2 g_{ab} \quad (9)$$

and

$$N_{ab} = \frac{1}{12} e^{-4\phi} [3H_{a\lambda k} H_b{}^{\lambda k} - \frac{1}{2} g_{ab} H^2]. \quad (10)$$

Considering the axionic antisymmetric tensor, $B_{\nu\lambda}$, as an independent variable, the equation of motion for the axion is:

$$\nabla_\mu [e^{-4\phi} H^{\mu\nu\lambda}] = 0 \quad (11)$$

which can be simplified into a wave equation of motion for a massless scalar field, $h(t, r, z, \varphi)$, evolving coupled to the dilaton [14]

$$\square h + 4\nabla^\mu \phi \nabla_\mu h = 0 \quad (12)$$

by writing

$$e^{-4\phi} H^{\mu\nu\lambda} = \epsilon^{\mu\nu\lambda k} h_{,k} \quad (13)$$

with $\epsilon^{\mu\nu\lambda k}$ the antisymmetric Levi Civita tensor given by

$$\epsilon_{abcd} \epsilon^{abcf} = -3! \delta_d^f. \quad (14)$$

In terms of h , the energy-momentum tensor for the axion, (10), then becomes:

$$N_{ab} = \frac{1}{2} e^{4\phi} [h_{,a} h_{,b} - \frac{1}{2} g_{ab} h_{,\alpha} h_{,\beta} g^{\alpha\beta}]. \quad (15)$$

To include the self-gravity of the string we require a metric which exhibits the symmetries of the sources, namely the boost invariance for the matter and dilatonic sectors ($T_0^0 = T_z^z$, $S_0^0 = S_z^z$). Therefore we try first the static cylindrically symmetric metric given in (1). Consistency with the assumed symmetry then requires $N_{00} = -N_{zz}$, i.e., $h_{,t} = h_{,z} = 0$ and also $N_{R\varphi} = 0$, i.e., $h_{,R} = 0$ or $h_{,\varphi} = 0$. Therefore there are two possible *Ansätze* for the axion: either $h(R)$ or $h(\varphi)$. Consistency of the constraint equation (given later by (17c)) also requires N_R^R to be a function of R and therefore for the ansatz $h(\varphi)$ we require $h_{,\varphi} = h$, i.e., $h(\varphi) = h\varphi$, with h a constant.

(i) $h(R)$

Let us first assume $h(R)$. Integrating the equation of motion for the axion (12) one gets:

$$\alpha h' = h_0 e^{-4\phi} \quad (16)$$

where h_0 is an integrating constant, with the energy-momentum tensor for the axion given by: $\hat{N}_0^0 = \hat{N}_z^z = \hat{N}_\varphi^\varphi = -\hat{N}_R^R = \frac{1}{4} e^{-\gamma+4\phi} h'^2$ where the prime means now the derivative with respect to R .

The equations of motion for the geometry and for the dilaton are respectively:

$$\alpha'' = -\alpha e^\gamma V(\phi) - \epsilon \alpha e^\gamma (\mathcal{E} - \mathcal{P}_R) \quad (17a)$$

$$(\alpha\gamma')' = -\alpha e^\gamma V(\phi) + \epsilon \alpha e^\gamma (\mathcal{P}_R + \mathcal{P}_\varphi) \quad (17b)$$

$$\alpha'\gamma' = -\frac{1}{2} \alpha e^\gamma V(\phi) + \frac{\alpha\gamma'^2}{4} + \alpha\phi'^2 + \epsilon \alpha e^\gamma \mathcal{P}_R + \frac{h_0^2}{4\alpha} e^{-4\phi} \quad (17c)$$

$$(\alpha\phi')' = \frac{\alpha e^\gamma}{4} \frac{\partial V}{\partial \phi} + \epsilon(a+1) \alpha e^\gamma \mathcal{E} - \frac{1}{2} \epsilon \alpha e^\gamma (\mathcal{P}_R + \mathcal{P}_\varphi) + \frac{h_0^2}{2\alpha} e^{-4\phi} \quad (17d)$$

while the equations of motion for the vortex sector are given by

$$\frac{1}{\alpha} (\alpha X')' = \frac{X P^2}{\alpha^2} e^{2\gamma} + \frac{X}{2} (X^2 - 1) e^\gamma \quad (18a)$$

$$\alpha \left(\frac{P'}{\alpha} \right)' = -\gamma' P' + \frac{X^2 P}{\beta} e^\gamma \quad (18b)$$

-the same as for the dilatonic string, as the axion does not couple directly to matter. For further reference, the Bianchi identity now gives:

$$\begin{aligned} & \epsilon(\alpha e^\gamma \mathcal{P}_R)' - \epsilon \alpha' e^\gamma \mathcal{P}_\varphi - \frac{\epsilon}{2} \alpha \gamma' e^\gamma [\mathcal{P}_R - \mathcal{P}_\varphi - 2\mathcal{E}] \\ & + \alpha' \phi'^2 + (\alpha \phi'^2)' - \frac{\alpha}{2} e^\gamma \phi' \frac{\partial V}{\partial \phi} + \alpha e^{4\phi} h' [h' (\frac{\alpha'}{2\alpha} + \phi') + \frac{h''}{2}] = 0 \end{aligned} \quad (19)$$

Let us start to look for the string's gravitational effects at very far distances from its core, where for non-singular local strings we are in vacuum.

Consider first the massless dilaton case, $V(\phi) \equiv 0$ for which the equations of motion for the geometry and the dilaton become:

$$\alpha'' = 0 \quad (20a)$$

$$(\alpha \gamma')' = 0 \quad (20b)$$

$$\alpha' \gamma' = \frac{\alpha \gamma'^2}{4} + \alpha \phi'^2 + \frac{h_0^2}{4\alpha} e^{-4\phi} \quad (20c)$$

$$(\alpha \phi')' = \frac{h_0^2}{2\alpha} e^{-4\phi}. \quad (20d)$$

These equations are only consistent for the trivial case where the axion is “switched off” and so the results in [11] for the dilatonic string still hold. Indeed one gets:

$$\alpha = dR + b \quad (21a)$$

$$\gamma = C_3 + \frac{C_2}{d} \text{Ln} [dR + b] \quad (21b)$$

$$\phi = \text{Ln} \left[e^{-8\phi_0} \frac{8C}{(\text{Cosh}(A))^2} \right]^{-\frac{1}{4}} \quad (21c)$$

with

$$A = \frac{\sqrt{8C}}{d} \text{Ln} \left[\frac{dR + b}{C_1} \right] \quad (22)$$

where C , C_1 , C_2 , C_3 , d and b are integrating constants.

From the dilaton equation (20d) it comes that C is defined such that:

$$(\alpha \phi')^2 = -\frac{h_0^2}{4} e^{-4\phi} + 2C \geq 0 \quad (23)$$

and therefore $C > 0$. But now we find a contradiction:- integrating (17b) from the core and to lowest order in ϵ one gets $\alpha\gamma' = 0$, i.e., $C_2 = 0$. Using the constraint (20c), $dC_2 = \frac{1}{4}C_2^2 + 2C$, this gives $C = 0$ which contradicts the fact that $C > 0$. This means $h' = \phi' = 0$. Therefore the only vacuum solution is the trivial one where the axionic sector is “switched off” and the dilaton is constant. This result still holds in the string frame and is in agreement with those presented in reference [12] for massless dilatons in a spacetime of four dimensions.

Consider now massive dilatons ($V(\phi) = 2M^2\phi^2$, with M the dilaton mass). Apart from an intermediate annular region (bounded by the Compton wavelength of the dilaton), the long-range structure of the fields is as for the Einstein case because for distances from the core greater than the Compton wavelength of the dilaton the dilaton field is essentially fixed. Hence the equations of motion for the geometry and the dilaton become:

$$\alpha'' = -\alpha e^\gamma 2M^2\phi^2 \quad (24a)$$

$$(\alpha\gamma')' = -\alpha e^\gamma 2M^2\phi^2 \quad (24b)$$

$$\alpha'\gamma' = -\alpha e^\gamma M^2\phi^2 + \frac{\alpha\gamma'^2}{4} + \frac{h_0^2}{4\alpha}e^{-4\phi} \quad (24c)$$

$$0 = \alpha e^\gamma M^2\phi + \frac{h_0^2}{2\alpha}e^{-4\phi}. \quad (24d)$$

whose solution is:

$$\alpha - \alpha_0 = C_4 R \quad (25a)$$

$$\gamma - \gamma_0 = -2 \ln[\alpha_0 + C_4 R] \quad (25b)$$

$$h - h_1 = -2\phi_1 \frac{C_5}{C_4} \ln[\alpha_0 + C_4 R] \quad (25c)$$

$$\phi = \phi_1 \quad (25d)$$

with

$$-\frac{h_0^2 e^{-4\phi_1}}{2\phi_1} = C_5 = \alpha^2 e^\gamma M^2 \quad (26)$$

which from (24c) gives

$$\phi_1 = -\frac{1}{4} - \frac{1}{4} \sqrt{1 + 48 \frac{C_4^2}{C_5}} \quad (27)$$

where α_0 , C_4 , γ_0 , h_1 are integrating constants and C_5 is a positive constant.

The asymptotic solution for the axionic-dilatonic cosmic string is then, to lowest order:

$$ds^2 = e^{\gamma_0} [\alpha_0 + C_4 R]^{-2} [dt^2 - dR^2 - dz^2] - e^{-\gamma_0} [\alpha_0 + C_4 R]^4 d\varphi^2 \quad (28)$$

with $\phi = \phi_1$ and the axion having very strong asymptotic effects as given in (25c).

$$(ii) \quad h(\varphi) = h\varphi$$

Let us now consider the other ansatz for the axion, $h(\varphi) = h\varphi$. The axion is no more a dynamical field and so its equation of motion gives an identity. The elements of the energy-momentum tensor are: $\hat{N}_0^0 = \hat{N}_z^z = \hat{N}_R^R = -\hat{N}_\varphi^\varphi = \frac{1}{4\alpha^2}e^{4\phi}h^2e^\gamma$ and the axionic sector in the initial action (6) is given by $\frac{1}{12}e^{-4\phi}H_{\mu\nu\lambda}^2 = \frac{1}{2}e^{4\phi+\gamma}\frac{h^2}{\alpha^2}$.

The equations of motion for the geometry and the dilaton are now respectively:

$$\alpha'' = -\alpha e^\gamma V(\phi) - \epsilon \alpha e^\gamma (\mathcal{E} - \mathcal{P}_R) - \frac{h^2}{2\alpha} e^{2\gamma+4\phi} \quad (29a)$$

$$(\alpha\gamma')' = -\alpha e^\gamma V(\phi) + \epsilon \alpha e^\gamma (\mathcal{P}_R + \mathcal{P}_\varphi) \quad (29b)$$

$$\alpha'\gamma' = -\frac{1}{2}\alpha e^\gamma V(\phi) + \frac{\alpha\gamma'^2}{4} + \alpha\phi'^2 + \epsilon \alpha e^\gamma \mathcal{P}_R - \frac{h^2}{4\alpha} e^{2\gamma+4\phi} \quad (29c)$$

$$(\alpha\phi')' = \frac{\alpha e^\gamma}{4} \frac{\partial V}{\partial \phi} + \epsilon(a+1)\alpha e^\gamma \mathcal{E} - \frac{1}{2}\epsilon \alpha e^\gamma (\mathcal{P}_R + \mathcal{P}_\varphi) + \frac{h^2}{2\alpha} e^{2\gamma+4\phi} \quad (29d)$$

and again the equations of motion for the vortex fields are the same as for the dilatonic string. The Bianchi identity is now:

$$\begin{aligned} & \epsilon(\alpha e^\gamma \mathcal{P}_R)' - \epsilon \alpha' e^\gamma \mathcal{P}_\varphi - \frac{\epsilon}{2} \alpha \gamma' e^\gamma [\mathcal{P}_R - \mathcal{P}_\varphi - 2\mathcal{E}] \\ & + \alpha' \phi'^2 + (\alpha \phi'^2)' - \frac{\alpha}{2} e^\gamma \phi' \frac{\partial V}{\partial \phi} - \frac{h^2}{\alpha} e^{4\phi+2\gamma} \phi' = 0. \end{aligned} \quad (30)$$

Again, consider first the massless case, $V(\phi) \equiv 0$. Proceeding as for the previous ansatz we write the equations of motion for the geometry and the dilaton:

$$\alpha' = -\alpha\phi' + q \quad (31a)$$

$$\alpha\gamma' = p \quad (31b)$$

$$\alpha \frac{\alpha''}{2} + \alpha'(-2q - p) + (q^2 + \frac{1}{4}p^2) + \alpha'^2 = 0 \quad (31c)$$

$$(\alpha\phi')' = \frac{1}{2\alpha} h^2 e^{2\gamma+4\phi} \quad (31d)$$

at far distances from the core where it is vacuum, with the Bianchi identity giving

$$\alpha' \phi'^2 + (\alpha \phi'^2)' - \frac{h^2}{\alpha} e^{4\phi+2\gamma} \phi' = 0 \quad (32)$$

where p and q are integrating constants. In order to get the asymptotic solutions to those equations we write the constraint (31c) as the autonomous d.s.:

$$u' = -3u^2 + 2(2q + p)ut - 2(q^2 + \frac{p^2}{4})t^2 \quad (33a)$$

$$t' = -ut \quad (33b)$$

where $t = \frac{1}{\alpha}$ and $u = \frac{\alpha'}{\alpha}$. Writing $x = \frac{u}{t} = \alpha'$ one gets:

$$dx = \left(3x + \frac{2}{x}(q^2 + \frac{p^2}{4}) - 2(2q + p) \right) \frac{dt}{t} \quad (34)$$

which integrated gives

$$t(x) = C (Q(x))^{\frac{1}{6}} \text{Exp} \left[-\frac{\sqrt{2}(p+2q)}{3\sqrt{p^2-8pq+4q^2}} \arctan \left[\frac{\sqrt{2}(p+2q-3x)}{\sqrt{p^2-8pq+4q^2}} \right] \right] \quad (35)$$

with C a positive integrating constant and $Q(x) = p^2 + 4q^2 - 4px - 8qx + 6x^2$, a non-invertible expression for $\alpha(R)$ whose asymptotic regimes are then analysed.

When $|x| \rightarrow \infty$ the result from expression (35) is that $t \rightarrow \infty$ where

$$t(x) = \frac{4}{C_3^4} |x|^{\frac{1}{3}} \quad (36)$$

Therefore the solution is:

$$\alpha = \alpha_0 + C_3 R^{\frac{1}{4}} \quad (37a)$$

$$\gamma = \gamma_0 + p \left(\frac{4\alpha_0^2}{C_3^3} R^{\frac{1}{4}} - \frac{2\alpha_0}{C_3^2} \sqrt{R} + \frac{4}{3C_3} R^{\frac{3}{4}} - \frac{4\alpha_0^3}{C_3^4} \text{Ln} [\alpha_0 + C_3 R^{\frac{1}{4}}] \right) \quad (37b)$$

$$\phi = \phi_0 + q \frac{4\alpha_0^2}{C_3^3} R^{\frac{1}{4}} - q \frac{2\alpha_0}{C_3^2} \sqrt{R} + q \frac{4}{3C_3} R^{\frac{3}{4}} - (1 + q \frac{4\alpha_0^3}{C_3^4}) \text{Ln} [\alpha_0 + C_3 R^{\frac{1}{4}}] \quad (37c)$$

with C_3 a positive constant and α_0 , γ_0 and ϕ_0 being integrating constants. But now an inconsistency comes- consistency of equation (31d) requires $C_3 = h = 0$, i.e., the axionic sector for these solutions is “switched off”.

When $x \rightarrow x_{\pm}$, where $Q(x_{\pm}) = 0$, therefore from expression (35) $t \rightarrow 0$. As a result the solution is:

$$\alpha = \alpha_0 + x_{\pm} R \quad (38a)$$

$$\gamma = \gamma_0 + \frac{p}{x_{\pm}} \text{Ln} [\alpha_0 + x_{\pm} R] \quad (38b)$$

$$\phi = \phi_0 + \left(\frac{q}{x_{\pm}} - 1 \right) \text{Ln} [\alpha_0 + x_{\pm} R] \quad (38c)$$

which from (31d), as in previous asymptotic regimes, gives $h = 0$, i.e., the axionic sector for these solutions it is as well “switched off”.

Finally when $x \rightarrow 0$ then, from expression (35) $t \rightarrow t(0)$ and the solution is:

$$\alpha = \alpha_0 \quad (39a)$$

$$\gamma = \gamma_0 + \frac{p}{\alpha_0} R \quad (39b)$$

$$\phi = \phi_0 + \frac{q}{\alpha_0} R \quad (39c)$$

which, again, from (31d) gives $h = 0$. In conclusion the only vacuum solution is the trivial one where the axion is “*switched off*”. The same result is obtained in the string frame. Moreover no solution (in the Einstein frame) is regular at the origin where $\hat{N}_0^0 = \frac{1}{4\alpha^2} e^{4\phi+\gamma} h^2 \sim \frac{1}{R}$, i.e., $e^{4\phi} \sim R$ because from the dilaton equation (29d) one obtains $\phi \sim R$ whilst from the constraint (29c) one gets $\phi \sim \sqrt{R}$. Therefore solutions of the equations of motion are non-string type solutions.

It is worth examining the previous results. First we note that for the ansatz $h(\varphi) = h\varphi$ the elements of the energy-momentum tensor for the axion resemble those for a global string given by

$$\hat{T}_i^i = \frac{1}{R^2} [1, \quad 1, \quad 1, \quad -1] \quad (40)$$

and for which the energy per unit length is infinite, i.e., the string field has long-range gravitational interactions. This fact when combined with the absence of non-trivial vacuum solutions suggest that fields for the axion-dilaton string also have a long-range behaviour. Meanwhile taking a supermassive Higgs limit at far distances from the core of a string in Einstein gravity the boson vectorial mass vanishes and one is left with the Einstein-Maxwell equations which can be solved exactly. The Higgs boson becoming supermassive decouples from the electromagnetic field and the metric solution is the Melvin magnetic Universe given by [15]

$$ds^2 = (1 + \frac{1}{4} B^2 R^2)^2 [dt^2 - dR^2 - dz^2] - \frac{R^2}{(1 + \frac{1}{4} B^2 R^2)^2} d\varphi^2 \quad (41)$$

with B the strength of the magnetic field along the axis of symmetry. This suggests that in the presence of the axion and the dilaton one shall look for a supermassive Higgs limit and one shall obtain an exact metric solution - the dyonic Melvin universe which will be derived in the next section.

We also note that the asymptotic limit of the solution (41) is of the form:

$$ds^2 = \rho^4 [dt^2 - dz^2 - d\rho^2] - \frac{D^2}{\rho^2} d\varphi^2 \quad (42)$$

with D a positive constant which is classified as a “*global string*” in the sense of Comtet and Gibbons [16] - a static solution of the vacuum Einstein equations which is invariant under boosts along the direction of the string and which is invariant under translations and reflections in a plane orthogonal to the translations together with rotations about an axis of symmetry and reflections in planes through the axis, i.e., having “*whole cylinder* -

symmetry” and being non asymptotically locally flat. Of course, the metric of a global vortex (generated by some field theory) does not have this form [17] but the form

$$ds^2 = b_0(r - r_0)^2 \left[dt^2 - \cosh^2 \sqrt{b_0} t \, dz^2 \right] - dr^2 - C_0^2 d\varphi^2 \quad (43)$$

with b_0 and C_0 two constants ($b_0 > 0$) which is clearly non-static (if we force the metric to be static it becomes singular [18]) and has an event horizon at $r = r_0$. Also note that in the presence of a dilaton the asymptotic form of a global vortex is also non-static as was shown in [19].

3 Global string solutions: dyonic universes

Dilatonic Melvin Universe

The dilatonic Melvin magnetic universe is an exact solution of the Dilatonic-Einstein-Maxwell theory described by the action:

$$S = \int d^4x \sqrt{-g} \left[-R + 2(\nabla\phi)^2 - e^{-2a\phi} F^2 \right] \quad (44)$$

where F is the field strength of any $U(1)$ gauge field, A .

In the presence of a magnetic field B the equations of motion given by [20]:

$$\nabla_\mu \left[e^{-2a\phi} F^{\mu\nu} \right] = 0 \quad (45a)$$

$$\square\phi + \frac{1}{2} a e^{-2a\phi} F^2 = 0 \quad (45b)$$

$$R_{\mu\nu} = 2\nabla_\mu\phi\nabla_\nu\phi + 2e^{-2a\phi} F_{\mu\rho} F_\nu^\rho - \frac{1}{2} g_{\mu\nu} e^{-2a\phi} F^2. \quad (45c)$$

admit an exact solution, the dilatonic Melvin magnetic universe, given by [20]

$$ds^2 = \left(1 + \frac{a^2 + 1}{4} B^2 R^2 \right)^{\frac{2}{1+a^2}} \left[dt^2 - dR^2 - dz^2 \right] - \left(1 + \frac{a^2 + 1}{4} B^2 R^2 \right)^{-\frac{2}{1+a^2}} R^2 d\varphi^2 \quad (46a)$$

$$e^{-2a\phi} = \left(1 + \frac{a^2 + 1}{4} B^2 R^2 \right)^{\frac{2a^2}{1+a^2}} \quad (46b)$$

$$A_\varphi = -\frac{2}{(1 + a^2)B(1 + \frac{a^2+1}{4} B^2 R^2)} \quad (46c)$$

$$B^z = -\frac{B}{\left(1 + \frac{a^2+1}{4} B^2 R^2 \right)^{\frac{2(2+a^2)}{1+a^2}}} \quad (46d)$$

The equations of motion are duality invariant as they admit an $SL(2, R)$ electric-magnetic duality defined by:

$$\phi' = -\phi \quad (47a)$$

$$F'_{\mu\nu} = \frac{1}{2}e^{-2a\phi}\epsilon_{\mu\nu\rho\sigma}F^{\rho\sigma} \quad (47b)$$

which can be used to generate an electromagnetic Melvin universe and that will be generalised in the presence of an axion in the next section.

Dyonic Melvin Universe

In the presence of an axion, the Melvin magnetic universe can be generated after applying a Peccei-Quinn and a duality transformation to the dilatonic Melvin magnetic universe. Our starting point comes from the bosonic part of the four-dimensional effective action of low energy string theory that includes the terms [21] :

$$S = \int d^4x \sqrt{-g} \left[-R + 2(\nabla\phi)^2 + \frac{1}{12}e^{-4\phi}H_{\mu\nu\lambda}^2 - e^{-2\phi}F^2 \right] \quad (48)$$

where $H_{\mu\nu\lambda}$ describes the axion, with now

$$H = \partial B + \frac{1}{4}A \wedge F. \quad (49)$$

If we write the axion as in (13) the equations of motion can equally be derived from the action [21] :

$$S = \int d^4x \sqrt{-g} \left[-R + 2(\nabla\phi)^2 + \frac{1}{2}e^{4\phi}(\nabla h)^2 - e^{-2\phi}F^2 - h F \tilde{F} \right] \quad (50)$$

with $\tilde{F}$ the dual of F .

It is also convenient to combine the axion, h , and the dilaton, ϕ , in a complex scalar field, $\lambda = \lambda_1 + i\lambda_2 = h + ie^{-2\phi}$ called the “*axidilaton*”. In terms of λ the action simplifies to [21] :

$$S = \int d^4x \sqrt{-g} \left[-R + \frac{|\nabla\lambda|^2}{2\lambda_2^2} + \frac{i}{4}(\lambda F_+^2 - \bar{\lambda} F_-^2) \right] \quad (51)$$

where $F_{\pm} = F \pm i\tilde{F}$. The kinetic term for λ in (51) is invariant under the Peccei-Quinn shift of the axion, $h \rightarrow h + f$ and under the duality transformation $\lambda \rightarrow -\frac{1}{\lambda}$. The latter, for $h = 0$, reduces to the $\phi \rightarrow -\phi$ transformation of (47a).

The action as a whole is not invariant under this duality, because the Maxwell terms change sign, therefore the equations of motion given by [21]:

$$\frac{\nabla_\mu \partial^\mu \lambda}{\lambda_2^2} + i \frac{\partial_\mu \lambda \partial^\mu \lambda}{\lambda_2^3} - \frac{i}{2} F_-^2 = 0 \quad (52a)$$

$$\nabla_\alpha [\lambda F_+^{\alpha\beta} - \bar{\lambda} F_-^{\alpha\beta}] = 0 \quad (52b)$$

$$R_{\mu\nu} = \frac{\partial_\mu \bar{\lambda} \partial_\nu \lambda + \partial_\nu \bar{\lambda} \partial_\mu \lambda}{4\lambda_2^2} + 2\lambda_2 F_{\mu\rho} F_\nu{}^\rho - \frac{1}{2} \lambda_2 g_{\mu\nu} F^2 \quad (52c)$$

are not invariant either. In fact the Einstein equation (52c) is not invariant in general under the duality transformation [21]

$$\phi' = -\phi \quad (53a)$$

$$F'_+ = -\lambda F_+ \quad (53b)$$

$$F'_- = -\bar{\lambda} F_- \quad (53c)$$

as the second and third terms are non-invariant [21] and transform into [21]:

$$2\lambda_2 F_{\mu\rho} F_\nu{}^\rho - \frac{1}{2} \lambda_2 g_{\mu\nu} F^2 - \frac{\lambda_1 \lambda_2^2}{|\lambda|^2} (2F_{\mu\rho} \tilde{F}_\nu{}^\rho + 2F_{\nu\rho} \tilde{F}_\mu{}^\rho - g_{\mu\nu} F \tilde{F}) \quad (54)$$

Only if the extra terms in this expression vanish will the duality transformation map solutions into solutions. A duality transformation of the offending terms above shows that they transform into themselves with

$$2F_{\mu\rho} \tilde{F}_\nu{}^\rho + 2F_{\nu\rho} \tilde{F}_\mu{}^\rho - g_{\mu\nu} F \tilde{F} \rightarrow (\lambda_1^2 - \lambda_2^2) (2F_{\mu\rho} \tilde{F}_\nu{}^\rho + 2F_{\nu\rho} \tilde{F}_\mu{}^\rho - g_{\mu\nu} F \tilde{F}) \quad (55)$$

and therefore the Einstein equation is invariant when [21]

$$2F_{\mu\rho} \tilde{F}_\nu{}^\rho + 2F_{\nu\rho} \tilde{F}_\mu{}^\rho - g_{\mu\nu} F \tilde{F} = 0 \quad (56)$$

which implies that if this expression vanishes for a particular solution, it will also vanish for its duality transformation of that solution. Thus, beginning with a solution one may build up a family of solutions. Indeed the Melvin magnetic universe verifies the restriction (56) (the only non-vanishing field strength is $F_{R\varphi} = \frac{RB}{(1+\frac{B^2 R^2}{2})^2}$) and therefore one can generate a axionic-dilatonic Melvin electromagnetic universe. To do this we start from the dilatonic Melvin magnetic solution (46a) - (46b) and obtain an axion from a shift f on a Peccei-Quinn transformation, $\lambda' = \lambda + f$, then dualise it, $\lambda' = -\frac{1}{\lambda}$, and finally rescale,

$\lambda' = \lambda(f^2 + 1)$ [21]. One gets [21] :

$$\lambda' = -\frac{1}{\lambda + f} [f^2 + 1] \quad (57a)$$

$$h' = -\frac{f}{f^2 + e^{-4\phi}} [f^2 + 1] \quad (57b)$$

$$e^{-2\phi'} = \frac{e^{-2\phi}}{f^2 + e^{-4\phi}} [f^2 + 1] \quad (57c)$$

$$F' = \frac{1}{\sqrt{f^2 + 1}} [-fF + e^{-2\phi}\tilde{F}] \quad (57d)$$

$$A'_\varphi = -\frac{f}{\sqrt{f^2 + 1}} A_\varphi \quad (57e)$$

with A'_z , A'_t and A'_R unknown. For the dilatonic Melvin magnetic universe $F_{R\varphi} = \frac{RB}{k^2}$ and therefore:

$$F'_{tz} = \frac{e^{-2\phi}}{k^2 R} B_1 \quad (58a)$$

$$F'_{R\varphi} = \frac{R}{k^2} B_2 \quad (58b)$$

with

$$B_1 = \frac{1}{\sqrt{f^2 + 1}} B \quad (59a)$$

$$B_2 = -\frac{f}{\sqrt{f^2 + 1}} B \quad (59b)$$

where B_2 is the magnetic field of the axionic sector and $B^2 = B_1^2 + B_2^2$ with $k = 1 + \frac{B^2 R^2}{2}$.

Using (46b) with $a = 1$ to rewrite the dilaton before the transformation one finally gets the exact dyonic Melvin electromagnetic universe, with the axion and dilaton fields given by:

$$h(B_1, B_2, R) = \frac{B_2 B_1}{B_1^2 + B_2^2} \frac{1}{\left(1 + \frac{1}{4} R^4 (B_1^2 + B_2^2) B_1^2 + R^2 B_1^2\right)} \quad (60a)$$

$$e^{-2\phi(B_1, B_2, R)} = \frac{B_1^2}{B_1^2 + B_2^2} \frac{1 + \frac{1}{2} (B_1^2 + B_2^2) R^2}{\left(1 + \frac{1}{4} R^4 (B_1^2 + B_2^2) B_1^2 + R^2 B_1^2\right)} \quad (60b)$$

with the metric given by:

$$ds^2 = k \left[dt^2 - dR^2 - dz^2 \right] - \frac{1}{k} R^2 d\varphi^2. \quad (61)$$

As expected this solution is consistent with the dilatonic Melvin magnetic universe, (46a) - (46b), which corresponds to the case where there is no axion ($f = 0$) and so $B_1 = B$ ($B_2 = 0$) with $e^{-2\phi'} \equiv e^{2\phi} = \frac{1}{1 + \frac{1}{2} B^2 R^2}$.

Asymptotic global string solutions

At very far distances from the core the field solutions are then obtained from (60a)-(60b) by taking the limit $R \rightarrow \infty$. Therefore the axion and the dilaton are given by

$$h(B_1, B_2, R) \rightarrow \frac{4B_2}{R^4 B^4 B_1} \quad (62a)$$

$$\phi(B_1, B_2, R) \rightarrow \ln \left[\frac{B R}{\sqrt{2}} \right] \quad (62b)$$

with the metric for the axionic-dilatonic global string given by

$$ds^2 = \left(\frac{BR}{\sqrt{2}} \right)^2 \left[dt^2 - dR^2 - dz^2 \right] - \frac{2}{B^2} d\varphi^2. \quad (63)$$

4 Conclusions

In the presence of an axion we showed that the gravitational field of the string is quite different from the Einstein and dilatonic ones [11]. In fact the long range effects of the dilaton and of the axion exclude the existence of local string solutions. Instead we looked for far field solutions and presented expressions for the asymptotical fields which approaches the dilatonic Melvin magnetic universe in the presence of an axion. The axion is strongly damped to zero while the dilaton has very strong asymptotic effects similar to those for the dilatonic string for massless dilaton.

Acknowledgements

The author would like to thank Dr. Ruth Gregory for reading this manuscript, for useful discussions and suggestions.

References

- [1] A.Vilenkin and E.P.S.Shellard, *Cosmic strings and other Topological Defects* (Cambridge Univ. Press, Cambridge, 1994).
- [2] M. Hindmarsh, *Cosmic Strings-Dead Again?* - *hep-ph/9806469*.
- [3] B. Linet, *Class. Quant.Grav.* 7 L75 (1990).
- [4] B. Linet, *Gen. Rel. Grav.* 17 1109 (1985).

- [5] C.M.Will, *Theory and Experiment in Gravitational Physics* (Cambridge Univ. Press, Cambridge, 1981).
- [6] P. A. M. Dirac, Nature 139 323 (1937).
P. A. M. Dirac, Proc. Roy. Soc. A165 199 (1938).
- [7] T. Kaluza, Sit. Z. Preuss. Akad. Wiss. 966 (1921).
O. Z. Klein, Phys. 37 895 (1926).
- [8] P. Jordan, Zeit. Phys. 157 112 (1959).
- [9] C. Gundlach and M. Ortiz, Phys. Rev. D42 2521 (1990).
A. Barros and C. Romero, J.Math.Phys. 36 5800 (1995).
L. Pimentel and A.N. Morales, Rev. Mex. Fis. 36 199 (1990).
- [10] T. Damour and A. Vilenkin, Phys. Rev. Lett. 78, 2288, (1997).
B. Boisseau and B. Linet, Class.Quant.Grav. 14 3063 (1997).
C. Charmousis B. Boisseau and B. Linet, Phys.Rev. D57 5150 (1998).
O. Dando and R. Gregory, Phys.Rev. D58 023502 (1998).
- [11] R. Gregory and C. Santos, Phys. Rev. D56 1194 (1997).
- [12] A. Dabholkar, G. Gibbons, J. Harvey and F. Ruiz Ruiz, Nucl. Phys. B340 33 (1990).
- [13] E.Fradkin, Phys. Lett. B158 316 (1985).
C.Callan, D.Friedan, E.Martinec and M.Perry, Nucl. Phys. B262 593 (1985).
C.Lovelace, Nucl. Phys. B273 413 (1985).
- [14] E. Copeland, A. Lahiri and D. Wands, Phys. Rev. D50 4868 (1994).
- [15] M. A. Melvin, Phys. Lett. 8 65 (1964).
- [16] A. Comtet and G. W. Gibbons, Nucl. Phys. B299 719 (1988).
- [17] R. Gregory, Phys. Rev. D54 4955 (1996)

- [18] A. G. Cohen and D. B. Kaplan, Phys. Lett. B215 67 (1988)
R. Gregory, Phys. Lett. B215 663 (1988)
G. Gibbons, M. Ortiz and F. Ruiz, Phys. Rev. D39 1546 (1989).
- [19] O. Dando and R. Gregory, Phys. Rev. D58 023502 (1998).
- [20] F. Dowker, J. Gauntlett, D. Kastor and J. Traschen, Phys. Rev. D49 2909 (1994).
- [21] A. Shapere, S. Trivedi and F. Wilczek, Mod. Phys. Lett.A6 2677 (1991).